

# The VHR data region-based classification possibilities in the framework of Control with Remote Sensing of European CAP.

A. Carleer<sup>a,\*</sup>, E. Wolff<sup>a</sup>

<sup>a</sup> Institut de Gestion de l'Environnement et d'Aménagement du Territoire, Université Libre de Bruxelles, cp 130/02, 50 av. F.D. Roosevelt, 1050 Brussels, Belgium – (acarleer, ewolff@ulb.ac.be)

**Abstract - In the European CAP (Common Agricultural Policy) framework, the European Commission imposes on member states to prevent agricultural subsidy irregularities, and recommends the control with remote sensing (CwRS). In the framework of remote sensing procedure, the European Commission, by the way of his Joint Research Centre, advises the use of very high spatial resolution (VHR) satellite data. These data are extraordinary from the point of view of the spatial resolution but the use of these kinds of data involves some problems in the traditional per-pixel classification like the salt and pepper effect, the poor spectral resolution of the VHR data and the difficulty in classifying land use. The region-based classification could solve these problems and allows the use of other features on top of spectral features in the classification process. This study present the possibility of the VHR data region-based classification to the classification of the agricultural and rural land cover in the framework of the remote sensing control of the European Union CAP.**

## 1. INTRODUCTION

Since 1993, the European Commission DG Agriculture has promoted the use of "Control with remote Sensing" (CwRS) as appropriate control system within the Common Agricultural Policy (CAP) to prevent agricultural subsidy irregularities (Astrand et al., 2004). Since 1999, the MARS (Monitoring Agriculture with Remote Sensing) project at the JRC (Joint Research Centre) centralizes the satellite images acquisition and since a few years, advises the use of very high spatial resolution (VHR) satellite data in the control process. These data are extraordinary from the point of view of the spatial resolution and the gap between the spatial resolution of aerial photographs and satellite images strongly decreased (Carleer and Wolff, 2004), they have the advantages of the satellite images: ability to cover large areas on a routine basis and their large archives. Moreover, these data fulfill the accuracy imposed by the Commission to digitalize the agricultural parcels for the LPIS (Land Parcel Identification System) and can be used for the CwRS at the same time. In the 2004 campaign, for the VHR satellite acquisition there were 71 VHR sites covering 50000 km<sup>2</sup>, for a total number of 221 sites through Europe (Astrand et al., 2004).

In spite of these advantages, the use of this kind of data involves some problems in the traditional per-pixel classification. With the spatial resolution refinement, the internal variability within homogenous land cover units increases (Woodcock and Stralher, 1987, Alpin et al., 1997, Thomas et al., 2003). The increased variability decreases the statistical separability of land cover classes in the spectral data

space and this decreased separability tends to reduce per pixel classification accuracies.

Another disadvantage with the very high spatial resolution satellite images is the relatively poor spectral resolution (Herold et al., 2003).

To overcome these problems, a region based procedure can be used (De Wit and Clevers, 2004, Carleer et al., 2005). The segmentation, before classification, produces regions which are more homogeneous in themselves than with nearby regions and represent discrete objects or areas in the image. Each image region then becomes a unit analysis and makes it possible to avoid much of the structural clutter.

The segmentation has other advantages; it allows measuring and using a number of features, on top of spectral features (Herold et al., 2002a, 2003, Thomas et al., 2003). These features can be the surface, the perimeter, the compactness (area/perimeter<sup>2</sup>), the degree and kind of texture (Johnsson, 1994). The segmentation is one of the only method which ensures to measure the morphological features (surface, perimeter, shape...) (Segl and Kaufmann, 2001) which may be especially useful when very high spatial resolution data are available (Jensen and Cowen, 1999). The use of these additional features could allow to compensate for the low spectral resolution of VHR satellite images (Guindon, 2000, Herold et al., 2002b) and to increase the classification accuracy for spectrally heterogeneous classes (Lillesand and Kiefer, 1994).

## 2. OBJECTIVE

The features calculated on the regions can be numerous and they all cannot be used. Using all of them not only makes computer system slow, but compromises the accuracy of the classification (Penaloza and Welch, 1996, Guindon, 2000). A selection of relevant features for each class is then essential to minimize the redundant signatures. This study present a feature selection procedure used in order to show which features (spectral, textural and morphological) are useful for which classes.

Moreover the segmentation technique used in this study makes it possible to segment the image on several levels. Then, this study also ensures to show in which segmentation level the classes are well identified and with which features.

## 3. STUDY AREA AND IMAGE DATA

The study area is situated at 33 km in the south-east of Brussels (Belgium) and covers a large rural area principally characterized by croplands and pasture lands, but also by roads, isolated buildings, rivers, water bodies and villages. The image data are ortho-rectified SPOT and Ikonos images, acquired on March 17, 2004 and June 7, 2004, respectively.

\* Corresponding author.

The data cover a surface of 74 km<sup>2</sup> and the spatial resolutions are 20 m for the MIR SPOT band, 10 m for the green, red and NIR SPOT bands, 4 m for the blue, green, red and NIR Ikonos bands and 1 m for the panchromatic Ikonos band.

#### 4. METHOD

##### 4.1 Segmentation

The segmentation technique used in this study is a "Region Growing" technique implemented in the eCognition software developed by Definiens Imaging (Definiens Imaging, 2004). The procedure starts at each point in the image with one-pixel objects and in numerous subsequent steps smaller image objects are merged into bigger ones, throughout a pair-wise clustering process. This segmentation technique makes it possible to segment the image on several levels. Each level is made up of the merger of the lower level regions. In this study, 12 different levels were carried out on the basis of two different sets of Ikonos bands (only PAN, and multispectral bands + PAN). Only the Ikonos bands were used, because we wanted to keep the maximum accuracy for the boundaries location.

##### 4.2 Legend

The land cover legend used in this study is a hierarchical legend with two main classes (Table 1). They subdivided into 17 land cover mainly based on land cover colors and vegetation types or crops (level 2). The three grey levels, the very reflective surface and the red surface are further divided according to intra-class land use (level 3): transportation area and building.

Table 1. Legend

Level 1	Level 2	Level 3	
vegetation	shadow on vegetation	11	
	shrub and tree	12	
	herb (permanent meadow or pasture)	13	
	sugar beet	14	
	potato	15	
	ensilage maize	16	
	winter wheat	17	
	winter barley	18	
	chicory	19	
	textile flax	20	
non-vegetation	shadow on non-vegetation	21	
	water	22	
	red surface	23	transportation area <b>301</b>
	very reflective surface	24	
	light grey	25	building <b>302</b>
	medium grey	26	
	dark grey	27	

##### 4.3 Features

Forty-seven features were calculated on each region of each level for each class. These features can be distributed in three categories: the spectral, textural and morphological features.

The spectral features contain the mean of the 9 spectral bands, the mean of Ikonos and SPOT NDVI, and 4 ratios of Ikonos bands. The textural features contain the Standard Deviation of all bands and NDVI, and five Haralick textural features calculated on the PAN and NIR Ikonos band, and on the NIR SPOT band (Homogeneity, Contrast, Dissimilarity, Entropy and Angular Second Moment) (Haralick et al, 1973). The morphological features contain the area, length, width, length/width, perimeter and compactness of the regions.

##### 4.4 Feature selection

In order to find the most suitable features for each class at each segmentation level, the public domain program "Multispec" was used. This program was designed for the analysis of multispectral and hyperspectral image data. The most suitable features are found by calculation of class separability based on the Bhattacharyya distance. The Bhattacharyya distance is defined as:

$$B = \left[ \mu_1 - \mu_2 \right]^T \left[ \frac{\Sigma_1 + \Sigma_2}{2} \right]^{-1} \left[ \mu_1 - \mu_2 \right] + \frac{1}{2} \ln \frac{\frac{1}{2} \left[ \Sigma_1 + \Sigma_2 \right]}{\sqrt{|\Sigma_1| |\Sigma_2|}}, \quad (1)$$

where  $\mu_i$  and  $\Sigma_i$  are the mean vector and the covariance matrix of class  $i$ , respectively.

The Bhattacharyya distance is a sum where the first part represents the mean difference component and the second part the covariance difference component. A general problem results from having no defined thresholds in terms of class separability. However, this measure can be used to assess separability of land cover classes and to prioritize features that contribute most to discrimination among the land cover classes of interest (Herold et al., 2003). The Bhattacharyya distance provides a separability score between each land cover class for a given set of features. This information can be used to identify the features that contribute the largest amount of separation of these classes. The Bhattacharyya distance measure is derived from training areas selected in each level for each class. These training areas were selected by an expert on each segmentation level, as it is commonly done.

The top 5 feature combinations for best average separability are calculated for a combination of four features, for each level of the legend, on each level of the segmentation.

#### 5. FEATURE SELECTION RESULTS

Before beginning the tests on the classes of the different legend levels, we noticed that some crop classes were not covered by vegetation at the two dates (sugar beet, potato, ensilage maize and chicory). Then, we decided to carry out the test with all the legend level 2 classes in order to see the distribution of the classes.

##### 5.1 Discrimination of all legend level 2 classes

The best average separabilities were found with the segmentation level 5 of the Ikonos multispectral and PAN bands. The most relevant features are the mean of the red, NIR and NDVI Ikonos bands and the standard deviation of the Ikonos NDVI.



## 6. CONCLUSION

All the features found in the different tests are not surprising because the features found correspond to the visual impression. But it is impossible to choose visually specific features among all the features available, and a quantitative selection, like Bhattacharyya distance, is essential to do the choice. This study shows also the utility of the morphological and textural features to classify a lot of classes, principally the non-vegetation classes, in order to overcome the poor spectral resolution, and the segmentation utility in order to calculate these features on specific region without taking into account the nearby regions. The segmentation is one of the only methods which ensure to measure the morphological features (surface, perimeter, shape...) (Segl and Kaufmann, 2001). These morphological features are essential in the classification of the building and transportation area classes, when many studies used ancillary data (road buffer, D.E.M., existing maps ...) to classify these classes.

But, this study shows also that the multi-temporal spectral features are more effective than the spatial features, calculated thanks to the segmentation (textural and morphological features), for the discrimination of the vegetation classes and above all for the crop classes. But, the segmentation allows avoiding the "salt and pepper" effect in the classification process for these classes.

In the framework of the European CAP, a good identification of the non-vegetation classes is very important because these objects (classes) are not eligible for the subsidies. Moreover, the region-based classification of the classes like transportation area, water, tree, building ... is important to have stable objects from one year to another. The segmentation of the very high resolution satellite images are also a good base for the digitalization of the crop field.

## 7. ACKNOWLEDGEMENTS

The data used were funded by the European Commission. This study was carried out in collaboration with the DGA of the Ministry of the Walloon Region (Belgium). Alexandre Carleer would like to thank the FRIA (Fonds pour la Formation à la Recherche dans l'Industrie et l'Agriculture) for its support.

## 8. REFERENCES

- Aplin P., P.M. Atkinson, P.J. Curran, Fine spatial resolution satellite sensors for the next decade, *International Journal of Remote Sensing*, **18 (18)**, 3873-3881, 1997.
- Astrand P.-J., Wirnhardt C., Biagini B., Weber M., Hellerman R., Controls with remote sensing of (CAP) arable and forage area-based subsidies – a yearly more than 700 images and 3 m euro affair..., *Proceedings of SPIE Sensors, Systems, and Next-Generation Satellites VIII*, Vol. 5570, eds. Meynart R., Neeck S.P., Shimoda H., 577-588.
- Barnsley M. J., S.T. Barr, Distinguishing urban land-use categories in fine spatial resolution land-cover data using a graph-based, structural pattern recognition system, *Computer environment and urban systems*, **21 (3/4)**, 209-225, 1997.
- Carleer A.P., E. Wolff, Exploitation of very high resolution satellite data for tree species identification, *Photogrammetric Engineering and Remote Sensing*, **70 (1)**, 135-140, 2004.
- Carleer A.P., Debeir O., Wolff E., Assessment of very high resolution satellite image segmentations, accepted in *Photogrammetric Engineering and Remote Sensing*, 2005.
- Definiens Imaging, eCognition Professional user guide, URL:<http://www.definiens-imaging.com/>, last consultation: March 15 2005.
- De Wit, A. J. W., Clevers, J.G.P.W., Efficiency and accuracy of per-field classification for operational crop mapping, *International journal of remote sensing*, **25(20)**, 4091-4112, 2004.
- Guindon B., Combining diverse spectral, spatial and contextual attributes in segment-based image classification, ASPRS Annual Conference, Washington DC, May 22-26, 2000.
- Haralick R.M., K. Shanmugam, I. Dinstein, Textural features for image classification, *IEEE Transactions on Systems, Man, and Cybernetics*, **sme-3 (6)**, 610-621, 1973.
- Herold M., J. Scepan, A. Müller, S. Günter, 2002a, Object-oriented mapping and analysis of urban land use/cover using IKONOS data, *Proceedings of 22<sup>nd</sup> EARSEL Symposium "Geoinformation for European-wide integration"*, Prague, June 2002.
- Herold M., M. Gardner, D.A. Roberts, 2002b, The spectral dimension in urban land cover mapping from high-resolution optical remote sensing data, *Proceedings of the 3<sup>rd</sup> Symposium on Remote Sensing of Urban Area*, Istanbul, Turkey, June 2002.
- Herold M., M.E. Gardner, D.A. Roberts, 2003, Spectral resolution requirement for mapping urban areas, *IEEE Transactions on Geoscience and Remote Sensing*, **41 (9)**, 1907-1919.
- Jensen J.R., D.C. Cowen, Remote sensing of urban/suburban infrastructure and socio-economics attributes, *Photogrammetric Engineering and Remote Sensing*, **65 (5)**, 611-622, 1999.
- Johnsson K., Segment-based land-use classification from SPOT satellite data, *Photogrammetric Engineering and Remote Sensing*, **60 (1)**, 47-53, 1994.
- Lillesand T.M. and R.W. Kiefer, *Remote Sensing and Image Interpretation*, pp. 750, John Wiley & Sons, Inc., New York, 1994.
- Penaloza M.A., R.M. Welch, Feature selection for classification of polar regions using a fuzzy expert system, *Remote Sensing of Environment*, **58**, 81-100, 1996.
- Segl K., H. Kaufmann, Detection of small objects from high-resolution panchromatic satellite imagery based on supervised image segmentation, *IEEE Transactions on geoscience and remote sensing*, **39 (9)**, 2080-2083, 2001.
- Thomas N., C. Hendrix, R.G. Congalton, A comparison of urban mapping methods using high-resolution digital imagery, *Photogrammetric Engineering and Remote Sensing*, **69 (9)**, 963-972, 2003.
- Woodcock C.E., A.H. Strahler, The factor of scale in remote sensing, *Remote Sensing of Environment*, **21**, 311-332, 1987.